



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

*Problem V.* — Assuming the same conditions as in problem IV, prove that in any two points of an elliptic orbit, the resistance will vary directly as the distance to the upper focus, and inversely as the fifth power of the square root of the distance to the lower focus.

*Problem VI.* — If the two forces are equal in circular orbits; and if  $a$  is equal to the semi-major axis of the earth's orbit; and  $v, g$  and  $r$  are respectively equal to the orbital velocity, intensity of the orbital force of gravity, and the etherial resistance of the earth at its mean distance from the sun; and if  $a' =$  the semi-major axis, and  $b' =$  the semi-minor axis of any elliptic orbit,—prove that one of the values of  $x$ , in the equation,

$$x^3 - 4a'x^2 + 4a'^2x - v^2g^2b'^2a'^2 \div r^2 = 0,$$

will be equal to the length of the radius vector to that point in the ellipse, where the accelerating force will be exactly balanced by the resisting force.

NOTE BY PROF. M. C. STEVENS.—In the article on Repetends published in Nos. 1 and 2 of Vol. I, it is stated, p. 25, lines 7 to 10, that “the operation may be very much abbreviated by taking advantage of the well-known property of repetends, that after one half of the figures are obtained, the second half may be found by subtracting each figure of the first half successively from 9;” and in demonstrating this property, on p. 27, it is assumed, if  $1 \div d$  be a fraction that reduces to a repetend, that in the reduction by division there will occur the remainder  $d - 1$ .

It should have been stated that this property applies only to such fractions reducing to pure repetends as have for denominators *prime numbers*.

It is therefore evident that the property is not applicable to the repetends resulting from  $\frac{1}{27}, \frac{1}{63}, \frac{1}{81}$ , &c.

The restriction is implied in the proof, since the remainder  $d - 1$  only occurs in case the denominator is a prime number.

NOTE BY THE EDITOR. — The solution of problem 105, given at p. 127, is defective, because the conclusion is virtually assumed by placing  $q - p = m - A + B - C + D - \&c.$  If we write instead,  $p - q = -m + A - B + C - D \&c.$ , we shall prove, in a similar manner, that  $p > q$ . We subjoin Dr. Nelson's solution of 105, which is entirely rigorous.

“The number of odd selections (1 at a time, 3 at a time, &c.) out of  $n$  shot

$$= S = n + \frac{n(n-1)(n-2)}{3!} + \frac{n(n-1)(n-2)(n-3)(n-4)}{5!} + \&c.$$

The number of even selections (2 at a time, 4 at a time, &c.)

$$= S' = \frac{n(n-1)}{2!} + \frac{n(n-1)(n-2)(n-3)}{4!} + \&c. \quad \text{Now } (1-1)^n = 0$$

$$= 1 - n + \frac{n(n-1)}{2!} - \frac{n(n-1)(n-2)}{3!} + \frac{n(n-1)(n-2)(n-3)}{4!} - \&c.$$

The sum of the negative terms in this series  $= S$ , and the sum of the positive terms  $= S' + 1$ . Hence  $S = S' + 1$ , or  $S - S' = 1$ . That is, the possible odd selections exceed the possible even selections, and therefore the chances are in favor of drawing an odd number."

### SOLUTION OF PROBLEMS IN NUMBER FIVE.

SOLUTIONS of problems in number five have been received as follows:

From R. J. Adcock, 136; Marcus Baker, 125, 126, 127, 130 and 133; Henry Gunder, 125, 126, 127, 128, 130, 133 and 136; William Hoover, 125, 126, 130, 133 and 136; Henry Heaton, 125, 128 and 135; Christine Ladd, 129; Prof. H. T. J. Ludwick, 130, 135 and 136; Artemas Martin, 125, 133, 134 and 136; D. J. Mc. Adam, 125, 130 and 133; E. B. Seitz, 125, 127, 129, 130, 133, 134, 135 and 136; Prof. J. Scheffer, 125, 126, 127, 128, 129, 130 and 133.

125.—"From a point  $E$ , in a square field  $ABCD$ , lines drawn to the corners  $A$ ,  $B$  and  $C$  are found to be 50, 30 and 40 rods, respectively. Required the side of the field."

SOLUTION BY E. B. SEITZ, GREENVILLE, OHIO.

Construct the right angled isosceles triangle  $BEF$ , making  $BF = BE = b = 30$  rods; on  $EF$  construct the triangle  $AEF$ , making  $AE = a = 50$  rods, and  $AF = c = 40$  rods; and on  $AB$  construct the square  $ABCD$ , which is the square required; for we can easily prove that  $EC = AF$ .

Put  $\angle AEF = a$ . Then we have  $\cos a = (a^2 + 2b^2 - c^2) \div (2ab \sqrt{2})$ , and  $AB = \sqrt{[a^2 + b^2 - 2ab \cos(a + \frac{1}{4}\pi)]}$

$$= \frac{1}{2} \sqrt{[2a^2 + 2c^2 + 2 \sqrt{(4a^2b^2 + 4b^2c^2 + 2a^2c^2 - 4b^4 - a^4 - c^4)}]}$$

$$= 5 \sqrt{(82 + 6 \sqrt{119})} = 60.71496 \text{ rods.}$$

126.—"In a plane triangle are given the vertical angle  $A$  and the bisectors  $\beta$  and  $\gamma$  of the base angles  $B$  and  $C$ ; determine the triangle."